

## **BRICKER ILEOSTOMY**

## Home education

- 1. Personnel in the ward will explain the necessary care and home procedures before discharge
- Home-care with a stoma therapist will be arranged and telephone numbers will be issued.
- 3. You have to continue with stoma care yourself at home. Stock will be supplied to you
- 4. Stoma care is to be done as needed with bag and base changes. Aseptic techinique
- 5. Monitor colour and wound care of stoma, and report any abnormalities to your doctor
- 6. Maintain high fluid intake
- 7. Prevent constipation
- 8. Avoid lifting anything heavy
- 9. Avoid driving or being a passenger in a vechicle for 4 weeks
- 10. No sexual activities for approximately 4-6 weeks.

Read more overleaf...

- 11. Maintain a healthy diet
- 12. Wear your TED socks until your follow up appointment
- 13. Continue with anticoagulant medicaiton as prescribed
- 14. Complete medication prescribed by doctor
- 15. Encourage leg and chest exercises
- 16. Report any abnormalites in urine output such as blood/sediments
- 17. Protection and treatment of surrounding stoma skin is very important.

  Gention Violet can be used with red skin.
- 18. Utheral catheters can still be in stoma on discharge, monitor your urine output. Do not pull on catheters. Stitches will dissolve in 10-14 days and only then the catheter will come out. Don't be frightened when you see them in the stoma bag.
- 19. Leg bag can be attached to lleostomy bag for 1st week at home while recovering. Thereafter, leg bag should be removed and continue only with a stoma bag
- 20. lleostomy bag to be emptied correctly
- 21. Attend your follow-up appointment
- 22. Maintain daily exercises
- 23. Stop smoking.



Cnr Grosvenor & Pretorius St, Hatfield Pretoria, 0083, South Africa P.O Box 13271, Hatfield, 0028 Tel: (012) 423 4000 Fax: (012) 342 1233 Email: urology@urology.co.za

www.urology.co.za