

# **FEMALE-SLING**

### What to expect at home

- 1. You may have pain or discomfort in your vaginal area or legs for a few months
- 2. Light bleeding or discharge from the vagina is normal.

### Self-Care:

How to take care of your surgical incision (cut):

- 1. Keep a dry dressing over the incision. Change the dressing as needed (as instructed by your doctor.)
- You may shower after your surgery. Gently wash the incision with mild soap and rinse well. Gently pat dry. Showers are better than baths for a few weeks, until your incision has healed.

## Try to prevent constipation

- 1. Eat foods that have a lot of fibre
- Use stool softeners as prescribed by your doctor. Straining during bowel movements will put pressure on your incision
- 3. Drink extra fluids to help keep your stools loose.

#### Do's and Dont's

- 1. You may slowly start your normal household activities. Be careful not to get overtired
- 2. Walk up and down stairs slowly. Walk every day. Start slowly with 5 minute walks, 3 or 4 times a day. Slowly increase the length of your walks
- 3. Do not lift anything heavier than 2 litres of milk for at least 4-6 weeks. Lifting heavy objects puts too much stress on your incision
- 4. Do NOT do strenuous activities such as golfing, playing tennis, bowling, running, biking, weight lifting, gardening or mowing and vacuuming for 6-8 weeks. Ask your doctor when it is acceptable to start
- 5. You may start sexual activity after 6 weeks. Ask your doctor when it will be acceptable to start
- 6. Attend your follow-up appointment.

### When to Call the Doctor

Call your doctor or nurse if any of the following occurs:

- 1. Severe pain
- 2. Fever
- 3. Heavy vaginal bleeding
- 4. Vaginal discharge with an odour
- 5. A lot of blood in your urine
- 6. You have signs of infection (a burning sensation when you urinate, fever or chills).



Cnr Grosvenor & Pretorius St, Hatfield Pretoria, 0083, South Africa P.O Box 13271, Hatfield, 0028 Tel: (012) 423 4000 Fax: (012) 342 1233 Email: urology@urology.co.za